

## Assemblage composition and seasonal variations of waterbirds at Nav Talav, Amgaon in Gondia District of Maharashtra

Jha, Ashish Kumar

**Received:** August 29, 2017 | **Accepted:** October 21, 2017 | **Online:** December 31, 2017

### Abstract

Among various biological components of freshwater, the waterbirds play an important role not only in trophic dynamics of ecosystem but also in the indication of changes in the quality of water due to pollution or degradation because of their ability to respond quickly to such changes. Thirty waterbird species, mainly Anatidae (26.57%) and Ardeidae (20.50%), were recorded at Nav Talav, Amgaon in Gondia district of Maharashtra using monthly point counts. Little Egret, Red-crested Pochard and Cotton Pygmy-goose were common, while the globally-near threatened Black headed Ibis and 15 species of local or regional conservation concern were also observed. Seasonal variation in waterbird assemblage structure reflected the arrival and/or passage of migrants, and few species were present between May and August. Further studies are

required to uncover the factors contributing to the observed seasonal variations in bird assemblage within Nav Talav, Amgaon.

**Keywords:** Waterbirds | Anatidae | Ardeidae | Seasonal variation

### Introduction

Wetland ecosystems harbor loads of biodiversity as it is being rich in nutritional value and productivity (Gibbs, 1993; Paracuellos, 2006) and are one of ecologically important conservation concern sites owing to its trophic dynamics. The waterbirds uses wetland habitats either throughout their life or during certain part of their life for nesting, breeding, feeding, sheltering and migration stopovers (Weller, 1999; Getzner, 2002). They are beneficial not only in trophic dynamics of ecosystem but also in the indication of changes in the quality of water due to pollution or degradation because of their ability to respond quickly to such changes. The spatial and seasonal distributions of waterbirds are influenced by conditions such as habitat types, climatic conditions, resource stability and immediate

#### For Correspondence:

Department of Bioscience and Biotechnology,  
Banasthali University, Banasthali, Rajasthan, India  
Email: [pracheta.25@gmail.com](mailto:pracheta.25@gmail.com);  
[sunanda.kulshrestha@gmail.com](mailto:sunanda.kulshrestha@gmail.com)

human impact. Different workers like, Osmatston (1922), Singh (1929), Ali (1932), Ghazi (1962), Kannon (1980), Mujumdar (1984), Davidar (1985), Newton *et al.*, (1986), Jhingram (1988), Ghosal (1995), Wadatkar and Kasambe (2002), Yardi *et al.*, (2004), Kulkarni *et al.*, (2005) and Harney (2014) have examined waterbird communities for annual variations in abundance and species composition from different freshwater bodies of India.

The present study was carried to investigate to overall assemblage composition and seasonal variations of waterbirds at Nav Talav, Amgaon in Gondia district of Maharashtra. The results would be useful in suggesting strategies for the conservation of Nav Talav and the waterbird diversity that it contains.

## Materials and Methods

### Study sites

Nav Talav at Amgaon in Gondia district is situated in eastern region of Maharashtra state at the geographical coordinate of 20°359'0"N latitude and 80°384'0"E longitude (Fig-1).

### Waterbird Survey

Waterbird surveys were carried out in Nav Talav from January 2013 to December 2013 on a monthly basis covering a complete wet season (June to September) and dry season (March to June); rainy days were avoided because rain interfere with visibility (Ralph *et al.*, 1993). Birds were identified by sight using binoculars (Olympus 10×50). During field studies, guidebooks were used to identify the birds (Ali, 2002, Grimmett *et al.*, 2011 and Manakadan *et al.*, 2011).

Obligate wetland users waterbirds from the avian fauna were selected and all other species were excluded from the analysis (Weller, 1999). All individuals of each waterbird species were counted in each visit within several parts of Nav Talav following the bird census techniques (Bibby *et al.*, 2000).

Since preliminary observations showed that waterbirds started to enter the study site for resting and foraging when the water level was high, all waterbird surveys were conducted when the water level was high. Waterbirds were defined as comprising species of cormorants, herons, egrets, ibises, ducks, rails and jacanas normally associated with wetlands and also birds species in the families Charadriidae (Lapwing) and Scolopacidae (Sandpipers) in the present study.

**The relative diversity (RDi)** of families was calculated by using following formula (Koli, 2014):

$$\text{RDi} = (\text{No. of birds species in the family} / \text{Total no. of species}) * 100$$


## Results and Discussion

A total of 824 waterbirds in 30 species were recorded in Nav Talav at Amgaon in Gondia district during the study period. The species belong to 6 orders: Pelecaniformes, Ciconiiformes, Anseriformes, Gruiformes, Charadriiformes, and Coraciiformes (Fig. 2). Based on foraging behavior and habitat use the waterbird families were grouped as follows: 1) surface and aerial diving birds (Phalacrocoracidae, Alcedinidae, Meropidae), 2) wading birds (Ardeidae, Ciconiidae, Threskiornithidae, Recurvirostridae, Charadriidae, Scolopacidae), 3) ducks (Anatidae), and 4) marsh birds. Most were

Anatidae (26.57%), while Ardeidae (20.50%) ranked the second among families in terms of relative diversity (Table- 1). This agreed with the finding of Kumar (2006), who reported that, Ardeidae to be the most dominant family in Bharathpuzha river basin in Kerala and Surana *et al.*, (2007) according to them Anatidae to be the most dominant family in Chimdi lake Nepal.



**Fig. 1:** A satellite view of Nav Talav, Amgaon in Gondia district of Maharashtra



**Fig. 2:** Abundance (no./sq.m) of waterbirds Order at Nav Talav, Amgaon in Gondia district of Maharashtra (January, 2013 to December 2013): (expressed in %)

The most common species (making up > 7% of total counts over the study period) were Red-crested Pochard (*Netta rufina*), Cotton Pygmy-goose (*Nettapus coromandelianus*), Little Egret (*Egretta garzetta*) and Little Cormorant (*Phalacrocorax niger*) (Table - 3). 21 species are very common, 03 are common, 04 are uncommon, 01 are occasional and 01 were rare to Nav Talav (Table - 2). Black-

headed Ibis (*Threskiornis melanocephalus*), Blue-tailed Bee-eater (*Merops philippinus*), Marsh Sandpiper (*Tringa stagnatilis*) and Great Cormorant (*Phalacrocorax carbo*), all of these species were relatively rare (< 0.5% of total counts). Similar observation was also recorded from Shrungarbandh lake (Bhandarkar and Paliwal, 2014), Zaliya lake (Puri, 2015) and Bothalkasa lake (Puri and Virani, 2016) in Gondia District of Maharashtra.

Of the 30 waterbird species recorded during the present study, 50% are of local or regional conservation concern (Kulkarni 2005, Kulkarni *et. al.*, (2006 (a), 2006 (b), 2006 (c)), among which Black-headed Ibis (*Threskiornis melanocephalus*), the globally-near threatened species (NT; Bird Life International, 2006), had the highest conservation value. Black-headed Ibis were recorded in Nav Talav in December and January, 2013 but the present study provides the first record of this species in Amgaon tehsil. Although only few individuals were recorded during the study period, 3 juveniles were seen roosting within the Nav Talav. These sighting records of Black-headed Ibis suggest that Nav Talav might be another location in Gondia District.

The Red-crested Pochard (*Netta rufina*) occupied first position in order of dominance in respect to total number of wetland birds (Table - 3). The population maximum of this species was in the month November. The second predominant species was Cotton Pygmy-goose (*Netta puscromandelianus*). This species exhibited its population peak in the month of November (Table-3). The third numerical dominant species was Little Egret

(*Egretta garzetta*). It showed its maxima in February (Table - 3). The other numerically important species observed were Spot-billed Duck (*Anas poecilorhyncha*), Tufted Duck (*Aythya fuligula*), Cattle Egret (*Bubulcus ibis*), Red-wattled Lapwing (*Vanellus indicus*), Asian Openbill (*Anastomus oscitans*) and Green Bee-eater (*Merops orientalis*). Great Cormorant (*Phalacrocorax carbo*), Black-headed Ibis (*Threskiornis melanocephalus*), Marsh Sandpiper (*Tringa stagnatilis*) and Blue-tailed Bee-eater (*Merops philippinus*) occurred in individual in very poor number in relation to total wetland bird community (Table - 3).

There was an obvious reduction in waterbird species richness in Nav Talav from March to September 2013 (when no more than 20 species were recorded in any month), while the number of species seen from October 2013 to February 2013 was greater (24 to 30). Nav Talav is an important stopover and wintering site for migrating birds, and the majority of birds in Gondia are either passage migrants or winter visitors. From mid-November to early January (winter migration), migrants birds pass through Gondia District from the northern breeding sites to wintering grounds near the equator and they fly back to the breeding sites, passing through and staying in Gondia district in April and May (Summer migration) (Viney *et al.*, 2005). All winter visitors and passage migrants leave Gondia district to their summer breeding grounds by the beginning of autumn (Dudgeon & Corlett, 2004). Since few of the waterbirds (20% of total) recorded in Nav Talav were either passage migrants or winter visitors, it is not surprising that the number of waterbird species recorded in Nav Talav dropped from

30 species in December to 19 species in February 2013, followed by a summer-low where only resident species including all egrets except the Intermediate Egret, and summer visitors were recorded.

Prey abundance, distribution and composition among areas have a direct influence on habitat use by waterbirds (Murkin & Kadlec, 1986; Colwell & Landrum, 1993; Sánchez *et al.*, 2006a; Sánchez *et al.*, 2006b). Aquatic invertebrates are an important food of wetlandbirds (Skagen & Oman, 1996; Pérez-Hurtado *et al.*, 1997; Gammonley & Laubhan, 2002), and positive correlations between waterbird and benthic invertebrates is common (Colwell & Landrum, 1993; Yates *et al.*, 1993; Safran *et al.*, 1997; Placyk & Harrington, 2004; Sánchez *et al.*, 2006a; Sánchez *et al.*, 2006b). Since the assemblage structures of aquatic macroinvertebrates in different areas have yet to be studied in Nav Talav, it is not possible to relate the distribution of wetland birds to their prey.

The greater counts of Little Egret, Red-crested Pochard and Cotton Pygmy-goose recorded in Nav Talav may be due to the fact that Nav Talav harbor loads of mollusca, crustacean and insects. The greater number of Cattle Egrets, which prey on insects, frogs and lizards (Strange, 2000), in Nav Talav may be explained by the presence of grazing cattle which helped to flush out preys (MacKilligan, 2005). Habitat selection by waterbirds has been shown to be affected by predation risk (Caldwell, 1986; Yasué, 2006). For instance, waders may avoid profitable feedings sites in the proximity of trees and shrub cover from which raptors launch surprise attacks (Cresswell & Whitfield, 1994).

Since the bird survey was conducted only once a month, it is inevitable that some waterbird species were missed, especially during the spring passage and autumn passage. In order to have a better understanding of the relationship between the distributions of prey and waterbirds in Nav Talav, further studies on the invertebrate

assemblage structure and sediment characteristics, which affects the type and abundance of invertebrates (Yates *et al.*, 1993; Bolduc & Afton, 2003; Sarkar *et al.*, 2005) and waterbirds (Grant, 1984; Mouritsen & Jensen, 1992; Granadeiro *et al.*, 2007) will be needed.

Anatide	26.58	Charadriidae	3.28
Ardeidae	20.51	Recurvirostridae	2.18
Threskiornithidae	2.67	Scolopacidae	0.49
Ciconiidae	2.79	Jacaniidae	2.55
Alcedinidae	13.23	Phalacrocoracidae	11.41
Meropidae	3.76	Rallidae	10.56

**Table 1:** Relative diversity (RDi) of various families at Nav Talav, Amgaon in Gondia district of Maharashtra (January, 2013 to December 2013)

S. No	Order	Family	Scientific name	Common name	Population IUCN status	Residential status
1	Anseriformes	Anatide	<i>Nettapus coromandelianus</i>	Cotton Pygmy-goose	LC	RVc
			<i>Netta rufina</i>	Red-crested Pochard	LC	WVc
			<i>Anas poecilorhyncha</i>	Spot-billed Duck	LC	RC
			<i>Aythya fuligula</i>	Tufted Duck	LC	RO
2	Ciconiformes	Ardeidae	<i>Bubulcus ibis</i>	Cattle Egret	LC	RVc
			<i>Ardeola grayii</i>	Indian Pond Heron	LC	RVc
			<i>Egretta garzetta</i>	Little Egret	LC	RVc
			<i>Ardea purpurea</i>	Purple Heron	LC	RVc
			<i>Mesophoyx intermedia</i>	Intermediate Egret	LC	RVc
		Threskiornithidae	<i>Threskiornis melanocephalus</i>	Black-headed Ibis (White Ibis)	NT	SVUc
			<i>Pseudibis papillosa</i>	Black Ibis	LC	RVc
3	Coraciiformes	Alcedinidae	<i>Anastomus oscitans</i>	Asian Openbill	LC	RVc
			<i>Alcedo atthis</i>	Common Kingfisher	LC	RVc
			<i>Ceryle rudis</i>	Pied Kingfisher	LC	RVc
		Meropidae	<i>Halcyon smyrnensis</i>	White-throated Kingfisher	LC	RVc
			<i>Merops orientalis</i>	Green Bee-eater	LC	RVc
<i>Merops philippinus</i>	Blue-tailed Bee-eater	LC	WVUc			
4	Charadriiformes	Charadriidae	<i>Vanellus indicus</i>	Red-wattled Lapwing	LC	RVc
		Recurvirostridae	<i>Himantopus himantopus</i>	Black winged Stilt	LC	RC
		Scolopacidae	<i>Tringa stagnatilis</i>	Marsh Sandpiper	LC	WVUc
			<i>Metopidius indicus</i>	Bronze-winged Jacana	LC	RC
			<i>Hydrophasianus chairurgus</i>	Pheasant-tailed Jacana	LC	RUc
5	Pelecaniformes	Phalacrocoracidae	<i>Phalacrocorax carbo</i>	Great Cormorant	LC	WVc
			<i>Phalacrocorax fuscicollis</i>	Indian Cormorant	LC	WVc
			<i>Phalacrocorax niger</i>	Little Cormorant	LC	RVc
6	Gruiiformes	Rallidae	<i>Porzana pusilla</i>	Baillon's Crake	LC	WVRr
			<i>Gallinula chloropus</i>	Common Moorhen	LC	RVc
			<i>Porphyrio porphyrio</i>	Purple Swampphen		RVc
			<i>Amaurornis phoenicurus</i>	White – breasted Waterhen	LC	RVc
			<i>Fulica atra</i>	Common Coot	LC	RVc

NT-Nearly Threatened, LC- Least Concern, R- Resident, WV- Winter Visitor, SV- Summer visitor Rr-Rare (<5%), O – Occasional (5-24%), Uc-Uncommon (25-49%), C- Common (50-74%), Vc- Very common (75-100%)

**Table 2:** Systematic List of Bird Species at Nav Talav, Amgaon in Gondia district of Maharashtra (January, 2013 to December 2013)

Species	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.
<i>Nettapus coromandelianus</i>	0.196	0.26	0.11	0.036	0.03	0.02	N.C.	N.C.	0.25	0.12	0.55	0.49
<i>Netta rufina</i>	0.52	0	0	0	0	0	N.C.	N.C.	0	0	0.64	0.203
<i>Anas poecilorhyncha</i>	0.08	0.07	0.12	0	0	0	N.C.	N.C.	0	0.4	0.39	0.345
<i>Aythya fuligula</i>	0.196	0	0	0	0	0	N.C.	N.C.	0	0	0.175	0.155
<i>Bubulcus ibis</i>	0.203	2.57	0.64	0.52	0.226	0.26	N.C.	N.C.	0.345	0.4	0.39	0.08
<i>Ardeola grayii</i>	0.47	0.04	0.003	0.04	0.01	0.04	N.C.	N.C.	0.035	0.045	0.04	0.04
<i>Egretta garzetta</i>	0.323	0.556	0.37	0.145	0.117		N.C.	N.C.	0.02	0.12	0.55	0.33
<i>Ardea purpurea</i>	0.053	0.016	0.023	0	0	0	N.C.	N.C.	0.045	0.004	0.035	0.056
<i>Mesophoyx intermedia</i>	0.023	0	0	0	0	0	N.C.	N.C.	0	0	0.11	0.115
<i>Threskiornis melanocephalus</i>	0	0	0	0	0	0	N.C.	N.C.	0	0	0.045	0.035
<i>Pseudibis papillosa</i>	0.06	0.006	0	0.01	0.004	0	N.C.	N.C.	0.12	0.004	0.04	0.12
<i>Anastomus oscitans</i>	0.07	0.22	0.086	0.018	0.03	0.6	N.C.	N.C.	0.5	0.04	0.19	0.155
<i>Alcedo atthis</i>	0.03	0.03	0.016	0.035	0.035	0	N.C.	N.C.	0.045	0.04	0.045	0.5
<i>Ceryle rudis</i>	0.023	0.02	0.013	0.005	0.02	0.11	N.C.	N.C.	0.02	0.03	0.175	0.115
<i>Halcyon smyrnensis</i>	0.07	1.04	1.71	0.213	0.18	0.05	N.C.	N.C.	0.23	0.4	0.59	0.35
<i>Merops orientalis</i>	0.153	0.14	0.043	0.026	0.017	0.13	N.C.	N.C.	0.125	0.045	0.08	0.065
<i>Merops philippinus</i>	0.047	0.04	0.003	0.04	0.01	0.04	N.C.	N.C.	0.005	0.035	0.04	0.04
<i>Vanellus indicus</i>	0.023	0.02	0.0133	0.005	0.003	0.05	N.C.	N.C.	0.29	0.05	0.01	0.25
<i>Himantopus himantopus</i>	0.06	0.006	0	0	0	0	N.C.	N.C.	0	0.03	0.04	0.12
<i>Tringa stagnatilis</i>	0.023	0	0	0	0	0	N.C.	N.C.	0	0	0.023	0.02
<i>Metopidius indicus</i>	0.12	0.03	0.03	0.015	0.003	0.05	N.C.	N.C.	0.29	0.05	0.01	0.025
<i>Hydrophasianus chairurgus</i>	0.023	0.02	0.013	0.005	0.02	0.11	N.C.	N.C.	0.02	0.005	0.035	0.04
<i>Phalacrocorax carbo</i>	0.005	0	0	0	0	0	N.C.	N.C.	0.013	0.01	0.004	0.007
<i>Phalacrocorax fuscicollis</i>	0.01	0.03	0	0	0	0	N.C.	N.C.	0.045	0.045	0.35	0.35
<i>Phalacrocorax niger</i>	0.23	0.12	0.18	0.08	0.03	0.145	N.C.	N.C.	0.117	0.12	0.55	0.33
<i>Porzana pusilla</i>	0.047	0.04	0.003	0.04	0.01	0.04	N.C.	N.C.	0.005	0.035	0.04	0.035
<i>Gallinula chloropus</i>	0.153	0.14	0.043	0.026	0.016	0.13	N.C.	N.C.	0.115	0.01	0.065	0.01
<i>Porphyrio porphyrio</i>	0.28	0.29	0.08	0.05	0.003	0.017	N.C.	N.C.	0.05	0.055	0.03	0.065
<i>Amaurornis sphenicurus</i>	0.07	0.22	0.086	0.018	0.03	0.6	N.C.	N.C.	0.5	0.04	0.1	0.125
<i>Fulica atra</i>	0.053	0.016	0.023	0.056	0.01	0.004	N.C.	N.C.	0.04	0.045	0.004	0.035

**Table 3:** Abundance of waterbirds obtained per month from all sampling sites at Nav Talav, Amgaon in Gondia district of Maharashtra (January, 2013 to December 2013): (expressed in % of total population)

## References

Ali, S. (1932): Flowers birds and birds flower in India. J. Bom. Nat. Hist. Soc. 35: 573-605.

Ali, S. (2002): The Book of Indian Birds, 13<sup>th</sup> revised Ed. Mumbai: Bombay Natural History Society.

Bibby, C. J.; Burgess, N. D.; Hill, D. A. and Mustoe, S. H. (2000): Birds census techniques. London: Academic Press, P. 302.

BirdLife International (2006): Platalea minor. 2007 IUCN Red List of Threatened Species. Available from <<http://www.iucnredlist.org/search/details.php/17485/summ>>.

Bhandarkar, S. V. and Paliwal, G. T. (2014): Biodiversity and conservation status of water birds in Shrungarbandh lake district Gondia, Maharashtra, India. Int. J. of Life Sciences, 2(3): 239-243.

- Bolduc, F. and Afton, A. D. (2003): Effects of structural marsh management and salinity on invertebrate prey of water birds in marsh ponds during winter on the Gulf Coast Chenier Plain. *Wetlands*, 23: 897-910.
- Caldwell, G. S. (1986): Predation as a selective force on foraging herons: effects of plumage color and flocking. *Auk*, 103: 494-505.
- Colwell, M. A. and Landrum, S. L. (1993): Nonrandom shorebird distribution and fine-scale variation in prey abundance. *Condor*, 95: 94-103.
- Cresswell, W. and Whitfield, D. P. (1994): The effects of raptor predation on wintering wader populations at the Tynninghame estuary, southeast Scotland. *Ibis*, 136: 223-232.
- Davidar, P. (1985): Ecological interactions between the mistletoes and their avian pollinators in south India. *J. Bom. Nat. Hist. Soc.*, 82: 45-60.
- Dudgeon, D. and Corlett, R. (2004): *The Ecology and Biodiversity of Hong Kong*. Friends of the Country Parks and Joint Publishing (HK) Company Ltd.
- Gammonley, J. H. and Laubhan, M. K. (2002): Patterns of food abundance for breeding waterbirds in the San Luis Valley of Colorado. *Wetlands*, 22: 499-508.
- Getzner, M. (2002): Investigating public decisions about protecting wetlands. *J. Environ. Manage.*, 64: 237-246.
- Ghazi, H. K. (1962): Piscivorous birds of Madras, Madras. *J. of fisheries*, 1(1): 106-107.
- Ghosal, D. N. (1995): Avifauna of conservation areas, No. 7, Fauna of Kanha Tiger Reserve. *Zoological survey of India (ZSI)*, 63-91.
- Gibbs, J. P. (1993): The importance of small wetlands for the persistence of local populations of wetland-associated animals. *Wetlands*, 13: 25-31.
- Granadeiro, J. P.; Santos, C. D.; Dias, M. P. and Palmeirim, J. M. (2007): Environmental factors drive habitat partitioning in birds feeding in intertidal flats: implications for conservation. *Hydrobiologia*, 587: 291-302.
- Grant, J. (1984): Sediment microtopography and shorebird foraging. *Marine Ecology Progress Series*, 19: 293-296.
- Grimmett, R.; Inskipp, C. and Inskipp, T. (2011): *Birds of the Indian Subcontinent*, 2<sup>nd</sup> Ed. London WCIB 3DP: Christopher Helm, 528.
- Harney, N. V. (2014): Avifaunal diversity of Ghotnimbala lake near Bhadrawati, Chandrapur, Maharashtra, India, *Int. J. of Life Science*, 2(1): 79-83
- Jhingran, V. G. (1988): *Fish and fisheries of India*. Hindustan Publishing cooperation, New Delhi.
- Kannan, P. (1980): Nector feeding adaptation of flower birds. *J. Bom. Nat. Hist. Soc.*, 75: 1036-1050.

- Koli, V. K. (2014): Diversity and status of avifauna in Todgarh-Raoli wildlife Sanctuary, Rajasthan, India. *J. of Asia-Pacific Biodiversity*, 7: 401-407.
- Kulkarni, A. N.; Kanwate, V. S. and Deshpande, V. D. (2005): Birds in and around Nanded city, Maharashtra. *Zoos Print Journal*, 20 (11), 2076-2078.
- Kulkarni, A. N.; Bhowte, C. S. and Kanwate V. S. (2006 a): Bird census in Nanded region Maharashtra. *Bioinfolet*, 3 (3): 173-178.
- Kulkarni, A. N.; Kanwate, V. S. and Deshpande V. D. (2006 b): Checklist of Birds of Shikhachiwadi Reservoir. District Nanded, Maharashtra. *J. Of Aqua. Biol*, 21(10): 80-85.
- Kulkarni, A. N. and Kanwate, V. S. (2006 c): Piscivorous birds of Dongarkheda irrigation tank. Dist Hingoli, Maharashtra. *Jr. Of Aqua. Biol.*, 21 (1), 86-87.
- Kumar, A. B. (2006): A checklist of Avifauna of the Bharthpuzha river basin, Kerala. *Zoo's Print*, 21(8): 2300-2355.
- MacKilligan, N. (2005): Herons, Egrets and Bitterns: Their Biology and Conservation in Australia. *CSIRO Publishing*, 133.
- Manakadan, R.; Daniel, J. C. and Bhopale, N. (2011): Birds of the Indian Subcontinent: A Field Guide. Mumbai: Bombay Natural History Society, 409.
- Mouritsen, K. N. and Jensen, K. T. (1992): Choice of microhabitat in tactile foraging dunlins *Calidris alpina*: the importance of sediment penetrability. *Marine Ecology Progress Series*, 85:1-8.
- Mujumdar, N. (1984): On a collection of birds from Bastar district, M.P. *Record Zoological survey of India*, Occasional paper No. 59:54.
- Murkin, H. R. and Kadlec, J. A. (1986): Relationships between waterfowl and macroinvertebrate densities in a northern prairie marsh. *Journal of Wildlife Management*, 50: 212-217.
- Newton, P. N.; Brudin, S. and Guy, J. (1986): The birds of Kanha Tiger Reserve Madhya Pradesh, India. *J. Bom. Nat. Hist. Society*, 83(3): 977-998.
- Osmaston, B. B. (1922): A Birds of Pachmari. *J. Bom. Nat. Hist. Soc.*, 28: 453-459.
- Paracuellos, M. (2006): How can habitat selection affect the use of a wetland complex by waterbirds? *Biodiversity and Conservation*, 15: 4569-4582.
- Pérez-Hurtado, A.; Goss-Custard, J. D. and García, F. (1997): The diet of wintering waders in Cádiz Bay, southwest Spain. *Bird Study*, 44: 45-52.
- Placyk, J. S. and Harrington, B. A. (2004): Prey abundance and habitat use by migratory shorebirds at coastal stopover sites in Connecticut. *Journal of Field Ornithology*, 75: 223-231.
- Puri, S. D. (2015): Avifaunal diversity of Malguzarilake at Zaliya near



- Amgaon in Gondia district (MS), India. *Int. J. of Life Sciences*, 3(3): 219-224.
- Puri, S. D. and Virani, R. S. (2016): Diversity and status of Avifauna from Bodalkasa lake in Gondia district, Maharashtra, India. *Int. J. of Life Sciences*, 4(2): 256-262
- Ralph, C. J.; Geupel, G.R.; Pyle, P.; Martin, T.E. and DeSante, D. F. (1993): *Handbook of Field Methods for Monitoring Landbirds*. Pacific Southwest Research Station, Albany, California, 41.
- Safran, R. J.; Isola, C. R.; Colwell, M. A. and Williams, O.E. (1997): Benthic invertebrates at foraging locations of nine waterbird species in managed wetlands of the northern San Joaquin Valley, California. *Wetlands*, 17: 407-415.
- Sánchez, M. I.; Green, A. J. and Castellanos, E.M. (2006a): Spatial and temporal fluctuations in presence and use of chironomid prey by shorebirds in the Odiel salt pans, south-west Spain. *Hydrobiologia*, 567: 329-340.
- Sánchez, M. I., Green, A. J. and Castellanos, E.M. (2006b): Temporal and spatial variation of an aquatic invertebrate community subjected to avian predation at the Odiel salt pans (SW Spain). *Archiv für Hydrobiologie*, 166: 199-223.
- Sarkar, S. K.; Bhattacharya, A.; Giri, S.; Bhattacharya, B.; Sarkar, D.; Nayak, D.C. and Chattopadhyaya, A. K. (2005): Spatiotemporal variation in benthic polychaetes (Annelida) and relationships with environmental variables in a tropical estuary. *Wetlands Ecology and Management*, 13: 55-67.
- Singh, T. C. N. (1929): A note of the pollination of *Erythring indica* by birds. *J. Bom. Nat. Hist.Soc.* 33: 960-462.
- Skagen, S. K. and Oman, H. D. (1996): Dietary flexibility of shorebirds in the Western Hemisphere. *Canadian Field-Naturalist*, 110: 419-432.
- Strange, M. (2000): *A Photographic Guide to the Birds of Southeast Asia: Including the Philippines and Borneo*. Periplus Editions (HK) Ltd.
- Surana, R.; Subba, B. R. and Limbu, K. P. (2007): Avian diversity during rehabilitation stage of Chimdi lake, Sunsari, Nepal. *Our Nature*, 5: 75-80.
- Viney, C.; Phillipps, K. and Lam, C. Y. (2005): *The Birds of Hong Kong and South China*, 8th edition. Information Services Department. Hong Kong Special Administrative Region, 255.
- Wadatkar, J. S. and Kasambe, R. (2002): Checklist of Birds from Pohara-Malkhed reserve forest, Distt. Amravati, Maharashtra. *Zoos. Print Journal*. 17(66): 807-811.
- Weller, M. W. (1999): *Wetland Birds: Habitat Resources and Conservation Implications*, Cambridge University Press, Cambridge.

- Yardi, D.; Patil, S. S. and Auti, R. G. (2004): Diversity of Avian Fauna from Salim Ali Lake of Aurangabad. Paper presented in 21st meet of birds lovers of Maharashtra held at Nanded on 3rd, 4th April - 2004.
- Yasué, M. (2006): Environmental factors and spatial scale influence shorebirds' responses to human disturbance. *Biological Conservation*, 128: 47-54.
- Yates, M. G.; Goss-Custard, J. D.; McGroarty, S.; Lakhani, K. H.; Durell, S. E. A.; Clarke, R. T.; Rispin, W. E.; Moy, I.; Yates, T.; Plant, R. A. and Frost, A. J. (1993): Sediment characteristics, invertebrate densities and shorebird densities on the inner banks of the Wash. *Journal of Applied Ecology*, 30: 599-614.